

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Left atrial appendage thrombi relate to easily accessible clinical parameters in patients undergoing atrial fibrillation transcatheter ablation: A multicenter study

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1632410> since 2019-06-01T16:47:01Z

Published version:

DOI:10.1016/j.ijcard.2017.04.014

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in INTERNATIONAL JOURNAL OF CARDIOLOGY, 241:218-222, 2017, 10.1016/j.ijcard.2017.04.014.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), 10.1016/j.ijcard.2017.04.014

The publisher's version is available at:

<http://linkinghub.elsevier.com/retrieve/pii/S0167527316328686>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/1632410>

**Left atrial appendage thrombi relate to easily accessible clinical parameters in patients
undergoing atrial fibrillation transcatheter ablation: a multicenter study**

Matteo Anselmino, MD PhD^a, Lucia Garberoglio, MD^a, Sebastiano Gili, MD^a, Emanuele Bertaglia, MD^c,
Giuseppe Stabile, MD^d, Raffaella Marazzi, MD^e, Sakis Themistoclakis, MD^f, Francesco Solimene, MD^g,
Simone Frea, MD^a, Walter Grosso Marra, MD^a, Mara Morello, MD^a, Marco Scaglione, MD^b, Roberto De
Ponti, MD, Prof^e, Fiorenzo Gaita, MD, Prof^a.

^a Division of Cardiology, Department of Medical Sciences, "Città della Salute e della Scienza" Hospital,
University of Turin, Italy;

^b Division of Cardiology, Department of Internal Medicine, Cardinal Massaia Hospital, Asti, Italy;

^c Department of Cardiac, Thoracic and Vascular Sciences, University of Padova, Padova, Italy

^d Clinica Mediterranea, Naples, Italy

^e Department of Heart and Vessels, Ospedale di Circolo e Fondazione Macchi, University of Insubria, Varese,
Italy.

^f Department of Cardiothoracic and Vascular Medicine, Dell'Angelo Hospital, Mestre-Venice, Italy

^g Clinica Montevergine, Mercogliano (AV), Italy

Word count: 3,948 words, 3 tables, 1 figure and 22 references

Short title: left atrial appendage thrombus and ablation

Corresponding author:

Sebastiano Gili, MD

Division of Cardiology, Department of Medical Sciences, "Città della Salute e della Scienza"
Hospital, University of Turin, Italy; Corso Bramante 88, 10126, Turin, Italy

Email : sebastiano.gili@gmail.com

Phone: +390116336022; Fax: +390116336015

1 **Abstract**

2 **Background.** Transesophageal echocardiography (TEE) is routinely performed before atrial
3 fibrillation (AF) transcatheter ablation to exclude the presence of left atrial (LA) or LA appendage
4 (LAA) thrombi. Aim of the study is to evaluate if easily accessible clinical parameters may relate to
5 the presence of LA or LAA thrombi to identify patients who could potentially avoid TEE.

6 **Methods and Results.** Between January 2012 and September 2014, data from 1,539 consecutive
7 patients undergoing TEE, as a work-up before AF transcatheter ablation, in six large volume centers
8 were collected. Baseline clinical features, CHA₂DS₂-VASc score, transthoracic echocardiography
9 and presence of thrombi at TEE were recorded. Exclusion criteria were valvular, hypertrophic or
10 dilated cardiomyopathy, previous heart surgery or an ejection fraction \leq 35%. Mean age was $59.6 \pm$
11 10.4 years, 1,215 (78.9%) were males; 951 (62.9%) presented in sinus rhythm (SR) on admission,
12 324 (21.1%) had undergone at least one previous ablation and 900 (58.5%) had CHA₂DS₂-VASc
13 score 0-1. Thrombi were encountered in 12 patients (0.8%). SR at TEE independently related to the
14 absence of thrombi (OR 5.15, 95% CI 1.38-19.02, p 0.015); in addition to this, no patient with a
15 CHA₂DS₂-VASc score 0-1 and SR on admission presented thrombi at TEE (specificity 100%,
16 p=0.011).

17 **Conclusion.** In a selected population of patients referred for AF ablation, LA/LAA thrombi
18 prevalence is low. No patients in SR with CHA₂DS₂-VASc score 0-1 presented LAA thrombi at TEE,
19 identifying a significant subset of patients who could potentially safely be spared from pre-procedural
20 TEE.

21

22 **Key words:** LAA thrombus; catheter ablation of atrial fibrillation; transesophageal echocardiography.

23

24

25

1 **Introduction**

2 Atrial fibrillation (AF) is the most common supraventricular arrhythmia. Its prevalence increases
3 with age and the number of patients is expected to increase in the next years¹. Percutaneous
4 transcatheter ablation has proven to be safe and effective and the number of procedures performed
5 worldwide is progressively increasing². A pre-procedure evaluation, using a transesophageal
6 echocardiogram (TEE), is commonly used to rule out the presence of thrombi in the left atrium (LA)
7 or left atrial appendage (LAA)^{2,3}. This examination is generally safe, technically simple and offers
8 the unique advantage to accurately rule out the presence of LA/LAA thrombi before a procedure
9 which can be associated with a significant risk of symptomatic³ and asymptomatic thromboembolic
10 events⁴. On the other side, this procedure can be time and resource consuming, often scarcely
11 tolerated by patients and presents a not negligible rate of complications, which can rarely be life-
12 threatening⁵. A very low prevalence of LA/LAA thrombi, moreover, is reported in patients
13 undergoing AF ablation⁶, making a discussion about the risk/benefit ratio of this exam in this setting
14 mandatory.

15 For these reasons, it would be of utmost importance to evaluate if any clinical feature may predict the
16 absence of LA/LAA thrombi and identify those patients who could potentially be spared from a TEE
17 before AF ablation. The present multicenter study assesses the prevalence of LA/LAA thrombi at
18 TEE in patients referred for AF ablation procedure in search of easily available clinical parameters
19 that may relate to LA/LAA thrombosis.

20
21
22
23
24
25

1 **Methods**

2 Medical records of all consecutive patients planned to undergo AF ablation from January 2012 to
3 September 2014 in six Italian high volume centers were retrospectively reviewed. Inclusion criteria
4 were all consecutive patients undergoing pre-procedural TEE before (Ö48 hours) elective AF
5 transcatheter ablation. In fact, at the time of collection of these data, all centres involved in the study
6 routinely performed TEE to all patients undergoing AF ablation. Exclusion criteria were the presence
7 of hypertrophic cardiomyopathy (HCM), dilated cardiomyopathy (DCM), restrictive cardiomyopathy
8 (RCM), constrictive pericarditis and valvular cardiomyopathy, defined as valvular defect of at least
9 moderate grade, previous heart surgery and a left ventricle ejection fraction (EF) Ö35%.

10 General and clinical data were recorded. Hypertension, dyslipidemia, disthyroidism, diabetes mellitus
11 were defined and classified according to the current international guidelines; CHA₂DS₂-VASc score
12 was computed for each patient.

13 AF was classified according to current European Society of Cardiology guidelines as paroxysmal,
14 persistent and long-standing persistent⁷. Arrhythmia duration was intended as time from the onset of
15 the first recorded arrhythmic episode.

16 Patients were treated according to the available latest AF ESC guidelines, particularly regarding
17 long-term oral anti-thrombotic prophylaxis⁷. Oral anticoagulation (OAC) prior to AF ablation could
18 be withdrawn or prosecuted according to physicians and treating centersøpreference; in any case, for
19 patients withdrawing OAC and/or not on long-term OAC a minimum of a 5 day-course of low-
20 molecular weight heparin at weight adjusted dose (80-100U/Kg every 12 hours) was prescribed.

21 All patients underwent pre-procedural trans-thoracic echocardiography, and LA/ LAA thrombi or
22 spontaneous echo contrast (SEC) were searched for by conventional TEE⁵. Thrombi were defined as
23 circumscribed and uniformly echodense intracavitary masses distinct from the underlying LA or
24 LAA endocardium⁸. Patients in whom a thrombus was described were temporarily withdrawn from
25 the AF ablation procedure. Ejection Fraction, antero-posterior LA diameter and LAA dimensions

were classified according to the current guidelines on Chamber Quantification ⁹.

The study was performed in accordance to the latest Declaration of Helsinki and patients at each center provided written informed consent to undergo TEE.

Statistical Analysis

Categorical variables are reported as counts and percentages, while continuous variables as means and standard deviations. Correlations between parameters and study groups were tested in cross tabulation tables by means of the Pearson Chi-Square or Fisher's Exact Test and by one-way ANOVA respectively for categorical and continuous variables. Sensitivity and specificity predictive values were assessed for clinical features reporting significant relation with LAA thrombi. Multivariate analysis was performed by a logistic binary regression model including parameters known to predict LAA thrombi and/or with a statistically significant relationship ($p < 0.05$) with the presence of LAA thrombi at univariate analysis.

A two sided p -value < 0.05 was considered statistically significant; all analyses were performed with SPSS 21.0 (IBM Corporation, Armonk, NY, USA).

1 **Results**

2 A total of 1,539 patients were enrolled, of whom 1,215 (78.9%) males; 864 (56.2%) presented
3 paroxysmal AF, 753 (48.9%) had hypertension and 66 (4.5%) suffered previous stroke or transient
4 ischemic attack (TIA); on admission, 951 (62.9%) were in sinus rhythm (SR, Table 1). Nine hundred
5 patients (58.5%) presented a CHA₂DS₂-VASc score of 0-1. Overall, 589 (56.2%) were on active
6 OAC treatment with a vitamin-K antagonists (VKA), 6 (0.4%) with novel oral anticoagulants
7 (NOAC). Overall, 164 (10.7%) patients were excluded due to the following exclusion criteria: 11 for
8 HCM, 39 for DCM, one for constrictive pericarditis, 79 for valvular heart disease, 18 for previous
9 heart surgery, 16 for EF ≤35%.

10 At TEE performed before the AF ablation procedure, LAA thrombi were detected in 12 patients
11 (0.8%). All patients with LAA thrombi were on OAC with VKA, excepted two with CHA₂DS₂-
12 VASc score 1. Among the patients with thrombi at TEE (Table 2), all of those with a CHA₂DS₂-
13 VASc score 0-1 were in AF at the moment of the TEE, while the remaining 3 patients that were in
14 SR during TEE presented a CHA₂DS₂-VASc score of 2 or higher. Seven of these patients were
15 undergoing a first ablation procedure; only one was in SR at the moment of TEE (CHA₂DS₂-VASc
16 score 3). At univariate analysis (Table 1), presence of AF during TEE and undergoing a redo
17 procedure, related to the presence of thrombi at TEE (p=0.024, p=0.001, respectively). No patient
18 with CHA₂DS₂-VASc score ≤1 and SR on admission presented thrombi at TEE (p=0.011, Figure 1).
19 Only one patient on SR undergoing a first procedure presented a thrombus at TEE (CHA₂DS₂-VASc
20 score 3). At multivariate analysis, adjusting for SR during TEE, CHA₂DS₂-VASc score 0-1,
21 undergoing a first procedure, only SR at TEE significantly related to the absence of thrombi (OR
22 5.15, 95% CI 1.38-19.02, p 0.015).

23 As expected, prevalence of SEC was significantly higher in patients with LAA thrombi (91
24 (6.1%) vs. 5 (50%), p < 0.001).

25 Specificity and predictive values for the absence of thrombi at TEE were assessed for the

clinical variables emerged at univariate analysis or known to relate to LA/LAA thrombi. Individually
CHA₂DS₂-VASc score as well as undergoing AF ablation for the first time reported a low specificity
(58.5% and 41.7, respectively), while a higher specificity (75.0%) was reported for heart rhythm on
admission and, even more, for presenting with SR at TEE while referred for the first procedure
(90.0%). No patients with CHA₂DS₂-VASc score 0-1 and SR at TEE presented LAA thrombi at TEE,
with a positive predictive value of 100% for the absence of thrombi for the combination of these two
parameters.

Discussion

The main findings of our study are 1) that prevalence of LA/LAA thrombi in selected, consecutive patients referred for AF ablation is extremely low; 2) presence of SR on admission and a low-intermediate CHA₂DS₂-VASc score, especially in patients undergoing first AF ablation, relate to the absence of LA/LAA thrombi at TEE.

AF ablation is an interventional procedure that is becoming a mainstay in the management of AF^{7,10}. Despite its widespread diffusion, questions concerning pre-procedural management, as the systematic need for TEE to exclude LA/LAA thrombi, still remain open. Not even the 2012 HRS/EHRA/ECAS consensus document on AF ablation reached an unanimous agreement on pre-procedural TEE indications, only suggesting TEE should be reserved to patients with AF on admission, especially when persisting from more than 48 hours¹¹. In fact, to date, given the difficulty of certifying OAC effect over the weeks preceding the procedure and the hazard of presenting a thrombi despite therapeutic OAC, the majority, but not all, of the centers routinely perform TEE before the AF ablation despite correct anticoagulation^{2,3,12}. In fact, in Europe, about 2 out of 3 patients undergo TEE prior to AF ablation, without major significant regional differences¹³.

Our study included a selected population of patients without major structural cardiomyopathies undergoing AF ablation and is, to the best of our knowledge, the largest available series assessing this issue. Despite the strict selection criteria, the population included in our study is well representative of a relevant share of subjects undergoing AF ablation in European centers in the time span considered for our analysis¹³. A very low prevalence of LA/LAA thrombi is reported, however the prospective study by Puwanant et al. reported a similar prevalence of LAA thrombi (0.6%)⁶; in the European Atrial Fibrillation Ablation Pilot Study 7 patients out of 1410 did not undergo AF ablation due to a LAA thrombus. A slightly higher prevalence was reported instead in other experiences (ranging between 1.6%-3.6%), that, however, included patients at higher risk of thrombotic events who were excluded in our multicenter study (e.g. history of congestive heart

failure, structural heart disease, cardiomyopathies and valvular heart disease)^{14, 15,16,17, 18,19}.

In the population included in the present study, SR at TEE, and referral for first AF ablation procedure significantly related to the absence of thrombi at TEE. Of note, also the presence of SEC at TEE related to the presence of LA/LAA thrombi (5.9 vs. 50.0%, respectively in patients without and with LA/LAA thrombi; $p<0.001$). Given the aim of the study to identify easily available features relating to the presence of LA/LAA thrombi, this parameter has intentionally not been considered within predictors; in any case, its presence, provided that no thrombi were detectable, did not prevent patients from undergoing the procedure.

SR at TEE was the only independent predictor of LA/LAA thrombi at multivariate analysis.

In addition, no patients in SR and with a CHA₂DS₂-VASc score of 0-1 presented thrombi at TEE.

These results are consistent with those reported in previous studies: in seven prior series, only 5 patients with a CHADS₂ or CHA₂DS₂-VASc score of 0 presented thrombi at TEE^{6,14,15,16,17,18,19}. Of these five patients only two were on SR at the time of TEE^{14,17,18,19}. The present findings therefore confirm that a clinical assessment combining two simple parameters, heart rhythm and CHA₂DS₂-VASc score, may help to identify a relevant amount of patients (35.1% in our series) with a very low probability of presenting LA/LAA thrombi at TEE prior to AF ablation. A history of a previous AF ablation, instead, related to the presence of LA/LAA thrombi also in a previous study of Wallace et al¹⁸. This result may depend on the fact that a previous transcatheter ablation may determine increased atrial fibrosis²⁰, while AF relapses following failure of a first procedure may relate to decreased atrial function and increased LA dimensions²¹, all predisposing substrates for thrombus formation and consequent thromboembolic risk.

In the present study, less than 60% of the patients not presenting a thrombus and slightly more than 30% of the patients in SR with low-intermediate CHA₂DS₂-VASc score were on long-term OAC before ablation. On the whole, all patients received at least a course of 5 days of low-molecular weight heparin before the TEE, remarking how the risk of LA/LAA thrombi is, in reality, in a general

population referred for AF ablation, quite low. These results may lead, in any case, to reconsider the appropriateness of executing TEE in all patients undergoing AF ablation.

In general, the present is a highly selected population with a low probability of LAA thrombi. Generalization of the present findings is therefore bound to subjects with similar characteristics. In fact, enrolment in the study was obtained after thorough TTE assessment in search of the declared exclusion criteria. Previous literature has proposed several TTE parameters, not considered in the present study, potentially related to an increased prevalence of LAA thrombi (i.e diastolic dysfunction, pulmonary hypertension, severe right atrial enlargement, poor LA function at speckle tracking²²). These TTE features, especially when the hereby-proposed features suggest sparing a TEE, therefore candidate as reliable parameters to minimize potential residual risks. Indeed TEE is a safe and low expensive exam, which offers undeniable results in the search of LA/LAA thrombi, but the risk-benefit ratio of its use in the AF ablation setting should be carefully re-assessed. TEE presents a low, but still not negligible, risk of complications (0.8-2.8%)⁵, and, in addition, may determine an unjustified increase in the health-care related costs other than being often poorly tolerated by the patients²³. In conclusion, in our opinion, a careful risk/benefit ratio evaluation and prospective studies are needed to corroborate the comparative safety of an extensive vs. selective use of TEE prior to AF ablation.

Study limitations

The following limitations must be taken into account. First, this report is retrospective: study cohort may therefore be biased by patient selection by physician in charge. Second, multivariate analysis is based on a low number of events and is consequently limited in statistical power. The results of our study have consequently to be considered primarily as hypothesis generating. Third, as the aim of our study was to identify easily available features relating to the presence of LA/LAA thrombi, advanced and TEE parameters were

intentionally not included in the analysis. More specifically, thorough diastolic (i.e. DTI E/e ϕ) or LA function (i.e. by strain and/or speckle tracking) assessment, number of LAA lobes and peak flow velocity were not available, as they were not part of clinical routine when excluding thrombi at pre-procedural TEE; eventually LAA morphology was not assessed due to need for 3D segmentation techniques not routinely performed in all the centers involved^{24 25}. Eventually, data concerning anticoagulation relate to prescribed and not effective treatment; for this reason, any conclusion on this topic has been avoided. Moreover, very few patients on NOAC were included, limiting generalization to patients on NOAC.

Conclusions

The present study shows that careful patient selection and a simple clinical assessment including CHA₂DS₂-VASc score, heart rhythm on admission, and ablation history may identify a relevant subset of patients with a very low probability of LA/LAA thrombi, who could be potentially safely spared from TEE before AF transcatheter ablation.

Acknowledgments: to all the cardiologists, cardiology fellows and paramedics assisting or performing the procedures and collecting the data that contributed to the realization of this study.

Conflicts of interest / Funding sources: none declared

- 1 **Author contributions**
- 2 Matteo Anselmino: Concept/design, Data analysis/interpretation, Drafting article, Statistics
- 3 Lucia Garberoglio: Concept/design, Data Collection, Critical revision of article
- 4 Sebastiano Gili: Concept/design, Data analysis/interpretation, Drafting article, Statistics
- 5 Emanuele Bertaglia: Concept/design, Critical revision of article, Approval of article
- 6 Giuseppe Stabile: Concept/design, Critical revision of article, Approval of article
- 7 Raffaella Marazzi: Critical revision of article, Data Collection
- 8 Sakis Themistoclakis: Concept/design, Critical revision of article, Approval of article
- 9 Franceso Solimene: Concept/design, Critical revision of article, Approval of article
- 10 Simone Frea: Critical revision of article, Data Collection
- 11 Walter Grosso Marra: Critical revision of article, Data Collection
- 12 Mara Morello: Critical revision of article, Data Collection
- 13 Marco Scaglione: Concept/design, Concept/design, Critical revision of article, Approval of article
- 14 Roberto De Ponti: Concept/design, Critical revision of article, Approval of article
- 15 Fiorenzo Gaita: Concept/design, Approval of article

16

17

18

19

20

21

22

23

24

25

26

27

1 References

-
- ¹ Go AS, Hylek EM, Phillips KA, Chang Y, Henault LE, Selby JV, Singer DE: Prevalence of diagnosed atrial fibrillation in adults: national implications for rhythm management and stroke prevention: the anticoagulation and risk factors in atrial Fibrillation (ATRIA) study. *JAMA* 2001; 285:2370-2375
- ² Cappato R, Calkins H, Chen SA, Davies W, Iesaka Y, Kalman J, Kim YH, Klein G, Natale A, Packer D, Skanes A, Ambrogi F, Biganzoli E: Updated Worldwide Survey on the Methods, Efficacy, and Safety of Catheter Ablation for Human Atrial Fibrillation. *Circ Arrhythm Electrophysiol* 2010; 3:32-38
- ³ Arbelo E, Brugada J, Hindricks G, Maggioni A, Tavazzi L, Vardas P, Anselme F, Inama G, Jais P, Kalarus Z, Kautzner J, Lewalter T, Mairesse G, Perez-Villacastin J, Riahi S, Taborsky M, Theodorakis G, Trines S; Atrial Fibrillation Ablation Pilot Study Investigators: ESC-EURObservational Research Programme: the Atrial Fibrillation Ablation Pilot Study, conducted by the European Heart Rhythm Association. *Europace*. 2012; 14(8):1094-103
- ⁴ Gaita F, Caponi D, Pianelli M, Scaglione M, Toso E, Cesarani F, Boffano C, Gandini G, Valentini MC, De Ponti R, Halimi F, Leclercq JF. Radiofrequency catheter ablation of atrial fibrillation: a cause of silent thromboembolism? Magnetic resonance imaging assessment of cerebral thromboembolism in patients undergoing ablation of atrial fibrillation. *Circulation*. 2010;122(17):1667-73.
- ⁵ Hahn RT, Abraham T, Adams MS, Bruce CJ, Glas KE, Lang RM, Reeves ST, Shanewise JS, Siu SC, Stewart

W, Picard MH: Guidelines for performing a comprehensive transesophageal echocardiographic examination: recommendations from the American Society of Echocardiography and the Society of Cardiovascular Anesthesiologists. J Am Soc Echocardiogr 2013; 26(9):921-64

⁶ Puwanant S, Varr BC, Shrestha K, Hussain SK, Tang WH, Gabriel RS, Wazni OM, Bhargava M, Saliba WI, Thomas JD, Lindsay BD, Klein AL: Role of the CHADS2 score in the evaluation of thromboembolic risk in patients with atrial fibrillation undergoing transesophageal echocardiography before pulmonary vein isolation. J Am Coll Cardiol 2009; 54(22):2032-9..

⁷ Camm AJ, Lip GY, De Caterina R, Savelieva I, Atar D, Hohnloser SH, Hindricks G, Kirchhof P: ESC Committee for Practice Guidelines (CPG). 2012 focused update of the ESC Guidelines for the management of atrial fibrillation: an update of the 2010 ESC Guidelines for the management of atrial fibrillation. Developed with the special contribution of the European Heart Rhythm Association. Eur Heart J 2012; 33(21):2719-47.

⁸ Aschenberg W, Schluter M, Kremer P, Schroder E, Siglow V, Bleifeld W. Transesophageal two-dimensional echocardiography for the detection of left atrial appendage thrombus. J Am Coll Cardiol 1986; 7(1):163-166.

⁹ Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernande L, Flachskampf FA, Foster E, Goldstein SA, Kuznetsova T, Lancellotti P, Muraru D, Picard MH, Rietzschel ER, Rudski L, Spencer KT, Tsang W, Voigt JU. Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the american society of echocardiography and the European association of cardiovascular imaging. Eur Heart J Cardiovasc Imaging. 2015;16(3):233-71

¹⁰ Anselmino M, D'Ascenzo F, Amoroso G, Ferraris F, Gaita F. History of transcatheter atrial fibrillation ablation. *J Cardiovasc Med (Hagerstown)* 2012; 13(1):1-8.

¹¹ Calkins H, Kuck KH, Cappato R, Brugada J, Camm AJ, Chen SA, Crijns HJ, Damiano RJ Jr, Davies DW, DiMarco J, Edgerton J, Ellenbogen K, Ezekowitz MD, Haines DE, Haissaguerre M, Hindricks G, Iesaka Y, Jackman W, Jalife J, Jais P, Kalman J, Keane D, Kim YH, Kirchhof P, Klein G, Kottkamp H, Kumagai K, Lindsay BD, Mansour M, Marchlinski FE, McCarthy PM, Mont JL, Morady F, Nademanee K, Nakagawa H, Natale A, Nattel S, Packer DL, Pappone C, Prystowsky E, Raviele A, Reddy V, Ruskin JN, Shemin RJ, Tsao HM, Wilber D: 2012 HRS/EHRA/ECAS Expert Consensus Statement on Catheter and Surgical Ablation of Atrial Fibrillation: recommendations for patient selection, procedural techniques, patient management and follow-up, definitions, endpoints, and research trial design. *Europace* 2012; 14(4):528-606

¹² Chen J, Todd DM, Hocini M, Larsen TB, Bongiorni MG, Blomström-Lundqvist C; Scientific Initiative Committee, European Heart Rhythm Association: Current periprocedural management of ablation for atrial fibrillation in Europe: results of the European Heart Rhythm Association survey. *Europace* 2014; 16(3):378-81

¹³ Riahi S, Arbelo E, Brugada J, Maggioni AP, Tavazzi L, Vardas P, Laroche C, Efremidis M, Xhaet O, Wo niak A, Van Gelder IC, Hjortshoj S. Regional differences in referral, procedures, and outcome after ablation for atrial fibrillation in Europe: a report from the Atrial Fibrillation Ablation Pilot Registry of the European Society of Cardiology. *Europace*. 2016;18(2):191-200

¹⁴ Scherr D, Dalal D, Chilukuri K, Dong J, Spragg D, Henrikson CA, Nazarian S, Cheng A, Berger

RD, Abraham TP, Calkins H, Marine JE: Incidence and predictors of left atrial thrombus prior to catheter ablation of atrial fibrillation. *J Cardiovasc Electrophysiol* 2009; 20(4):379-84

¹⁵ McCready JW, Nunn L, Lambiase PD, Ahsan SY, Segal OR, Rowland E, Lowe MD, Chow AW: Incidence of left atrial thrombus prior to atrial fibrillation ablation: is pre-procedural transoesophageal echocardiography mandatory? *Europace* 2010; 12(7):927-32.

¹⁶ Dorenkamp M, Sohns C, Vollmann D, Lüthje L, Seegers J, Wachter R, Puls M, Staab W, Lotz J, Zabel M: Detection of left atrial thrombus during routine diagnostic work-up prior to pulmonary vein isolation for atrial fibrillation: role of transesophageal echocardiography and multidetector computed tomography. *Int J Cardiol* 2013; 163(1):26-33

¹⁷ Han JH, Shin DH, Lee HJ, Kim YJ, Lee SH, Shim J, Uhm JS, Kim JY, Chang HJ, Pak HN, Lee MH, Joung B: Routine preprocedural transesophageal echocardiography might not be necessary for stroke prevention evaluation in AF patients on anticoagulation therapy. *Int J Cardiol* 2013; 168(3):1992-6.

¹⁸ Wallace TW, Atwater BD, Daubert JP, Voora D, Crowley AL, Bahnson TD, Hranitzky PM: Prevalence and clinical characteristics associated with left atrial appendage thrombus in fully anticoagulated patients undergoing catheter-directed atrial fibrillation ablation. *J Cardiovasc Electrophysiol* 2010; 21(8):849-52

¹⁹ Yamashita E, Takamatsu H, Tada H, Toide H, Okaniwa H, Takemura N, Sasaki T, Miki Y, Fuke E, Hayashi T, Sakamoto T, Nakamura K, Fukazawa R, Sato C, Goto K, Kaseno K, Kumagai K, Naito S, Hoshizaki H, Oshima S: Transesophageal echocardiography for thrombus screening prior to left atrial catheter ablation. *Circ*

²⁰ Badger TJ, Daccarett M, Akoum NW, Adjei-Poku YA, Burgon NS, Haslam TS, Kalvaitis S, Kuppahally S, Vergara G, McMullen L, Anderson PA, Kholmovski E, MacLeod RS, Marrouche NF: Evaluation of left atrial lesions after initial and repeat atrial fibrillation ablation: lessons learned from delayed-enhancement MRI in repeat ablation procedures. *Circ Arrhythm Electrophysiol* 2010; 3(3):249-59

²¹ Marsan NA, Tops LF, Holman ER, Van de Veire NR, Zeppenfeld K, Boersma E, van der Wall EE, Schalij MJ, Bax JJ: Comparison of left atrial volumes and function by real-time three-dimensional echocardiography in patients having catheter ablation for atrial fibrillation with persistence of sinus rhythm versus recurrent atrial fibrillation three months later. *Am J Cardiol* 2008; 102(7):847-53

²² Anselmino M, Gili S, Castagno D, Ferraris F, Matta M, Rovera C, Giustetto C, Gaita F. Do left atrial appendage morphology and function help predict thromboembolic risk in atrial fibrillation? *J Cardiovasc Med (Hagerstown)*. 2016 Mar;17(3):169-76.

²³ Cooper L, Candiotti K, Gallagher C, Grenier E, Arheart KL, Barron ME: A randomized, controlled trial on dexmedetomidine for providing adequate sedation and hemodynamic control for awake, diagnostic transesophageal echocardiography. *J Cardiothorac Vasc Anesth* 2011; 25(2):233-7.

²⁴ Anselmino M, Scaglione M, Di Biase L, Gili S, Santangeli P, Corsinovi L, Pianelli M, Cesarani F, Faletti R, Righi D, Natale A, Gaita F. Left atrial appendage morphology and silent cerebral ischemia in patients with atrial fibrillation. *Heart Rhythm*. 2014 Jan;11(1):2-7.

²⁵ Bai W, Chen Z, Tang H, Wang H, Cheng W, Rao L. Assessment of the left atrial appendage

structure and morphology: comparison of real-time three-dimensional transesophageal echocardiography and computed tomography. *Int J Cardiovasc Imaging*. 2016. doi: 10.1007/s10554-016-1044-4

Table 1. Baseline clinical and echocardiographic characteristics of the study population stratified according to the presence of left atrial appendage thrombi. Values reported as counts and percentage, if not differently stated. P value by Pearson Chi-Square/Fisher's Exact Test or ANOVA.

Characteristics	Overall population n = 1539	No LAA thrombus n = 1527	LAA thrombus n = 12	p value
Age (years)	59.6 ± 10.4	59.6 ± 11.3	59.1 ± 19.4	0.887
Male	1215 (78.9)	1206 (79.0)	9 (75%)	0.724
BMI	19.0 ± 12.7	19.0 ± 12.7	23.3 ± 11.5	0.412
Hypertension	753 (48.9)	745 (48.8)	8 (66.7)	0.256
Dislipidemia	237 (18.4)	235 (18.4)	2 (16.7)	0.990
Previous Stroke/TIA	66 (4.5)	65 (4.5)	1 (8.3)	0.800
Diabetes	89 (6.2)	89 (6.3)	0 (0)	0.371
CHA ₂ DS ₂ -VASc				
0	405 (26.3)	403 (26.4)	2 (16.7%)	0.297
1	495 (32.2)	492 (32.2)	3 (25.0%)	
2	382 (24.8)	379 (24.8)	3 (25.0%)	
≥ 3	257 (16.7)	253 (16.6)	4 (33.3%)	
Paroxysmal AF	864 (56.2)	860 (56.4)	4 (33.3)	
Persistent AF	620 (40.3)	615 (40.3)	5 (45.5)	
Long-standing persistent AF	48 (3.1)	48 (3.1)	0 (0.0)	
AF duration (months)	56.3 ± 66.0	56.4 ± 66.1	47.2 ± 48.5	0.756
First AF ablation procedure	1215 (79.0)	1208 (79.1)	7 (58.4)	<0.001
Sinus rhythm at admission	951 (62.9)	948 (63.2)	3 (25.0%)	0.024
OAC	589 (56.2)	579 (55.9)	10 (83.3%)	0.078
TT Echocardiographic parameters				
LVEF (%)	60.5 ± 6.6	60.0 ± 5.6	60.6 ± 6.5	0.819
LA volume (ml)	84.8 ± 39.3	117.7 ± 25.4	84.7 ± 39.4	0.149
Left atrium AP diameter (mm)	45.2 ± 6.7	45.2 ± 6.7	51.5 ± 8.9	0.061

BMI, body mass index; TIA, transient ischemic attack; AF, atrial fibrillation; AP, antero-posterior; LA, left atrium; LAA, left atrial appendage; LVEF, left ventricular ejection fraction; OAC, oral anticoagulation; TT, transthoracic

Table 2. Detailed clinical features of patients presenting with left atrial or left atrial appendage thrombus at transesophageal echocardiography.

Patient	Age	Sex	CHA ₂ DS ₂ -VASc score	First procedure	Long term OAC	Heart rhythm at TEE	Oral anticoagulation
1	67	M	2	NO	Yes	AF	Yes
2	65	F	5	No	Yes	SR	Yes
3	77	M	3	No	Yes	AF	Yes
4	63	M	1	Yes	No	AF	No
5	59	F	1	No	No	AF	No
6	65	M	2	No	Yes	SR	Yes
7	56	M	0	Yes	Yes	AF	Yes
8	63	M	3	Yes	Yes	SR	Yes
9	69	F	2	Yes	Yes	AF	Yes
10	58	M	2	Yes	Yes	AF	Yes
11	35	M	0	Yes	Yes	AF	Yes
12	67	M	1	Yes	Yes	AF	Yes

Table 3. Sensitivity and specificity predictive values for the clinical features showing a significant correlation for the absence of left atrial/left atrial appendage thrombi.

	Specificity	Sensitivity	PPV	NPV
CHA₂DS₂-VASc 0-1	58.5	58.3	99.4	1.1
SR at TEE	75.0	62.1	99.7	1.9
First Procedure	41.7	79.1	99.4	1.6
CHA₂DS₂-VASc score 0-1 and SR at TEE	100.0	35.2	100.0	1.2
SR at TEE and First Procedure	90.0	50.2	99.9	3.9
SR at TEE and CHA₂DS₂-VASc 0-1 and First Procedure	100.0	55.5	100.0	1.7

NPV, negative predictive value; PPV, positive predictive value

Figure titles and legends:

Figure 1. **Heart Rhythm, CHA₂DS₂-VASc score and Left Atrial or Left Atrial Appendage Thrombi.** Prevalence of left atrial or left atrial appendage thrombi stratified according to heart rhythm on admission and CHA₂DS₂-VASc score (SR, sinus rhythm; AF, atrial fibrillation).

